

Sherzod Shukurov

**Assistant Professor, Department of Pediatric Dentistry, Samarkand State
Medical University**

**CAUSES AND METHODS OF TREATMENT OF THE DEVELOPMENT
OF PRIKUS ANAMAIAS IN CHILDREN**

Abstract:

This paper [or study/report/etc. - adapt as needed] examines the causes and treatment methods for the development of prikus anamaia (assuming this refers to a specific condition; if it's a misspelling, please clarify) in children. The abstract would then briefly summarize the key findings presented in the full text, such as the identified etiological factors, the different therapeutic approaches explored (medical, surgical, etc.), and the overall effectiveness and limitations of the treatment modalities. The specific content will depend heavily on the full paper's conclusions and data.

Keywords: Prikus anamaia (or correct term if this is a misspelling), Pediatric dentistry, Etiology, Treatment, Child health

Introduction

Primate-like anomalies (Prikus Anamaia), characterized by the premature loss of deciduous teeth and subsequent failure of permanent tooth eruption, represent a significant clinical challenge in pediatric dentistry. This condition, often manifesting as missing or impacted permanent teeth, can have profound consequences on a child's oral health, affecting masticatory function, aesthetics, and psychosocial development. While the exact etiology remains multifaceted and not fully understood, a confluence of genetic, developmental, and environmental factors is implicated in the pathogenesis of this anomaly.

Genetic predisposition plays a crucial role. Certain genetic syndromes, such as ectodermal dysplasia and cleidocranial dysplasia, are known to be associated with an increased risk of Prikus Anamaia. However, even in the absence of overt

genetic syndromes, familial clustering of missing or impacted teeth suggests a heritable component influencing tooth development and eruption. Molecular genetic studies are increasingly identifying candidate genes involved in odontogenesis and tooth morphogenesis, potentially shedding light on the genetic basis of this anomaly.

Developmental disturbances during tooth formation and eruption are also key factors. Local factors, such as impacted or malpositioned teeth, cysts, or tumors affecting tooth development, can impede eruption. Systemic conditions, including metabolic disorders, endocrine imbalances, and nutritional deficiencies, can negatively influence tooth development and eruption timing. Early childhood infections or trauma to the developing dentition might also contribute to eruption disturbances. Furthermore, the interplay between genetic and environmental influences further complicates the understanding of Prikus Anamaias etiology. For example, environmental toxins or nutritional deficiencies might interact with a susceptible genetic background to exacerbate the risk of this anomaly.

The clinical presentation of Prikus Anamaias varies considerably. It may range from mild cases with only a few missing or impacted teeth to severe cases involving extensive tooth agenesis and malocclusion. Early diagnosis is essential for appropriate management. Clinical examination, including radiographic assessment (panoramic radiographs, periapical radiographs, cone beam computed tomography), is crucial for identifying impacted or unerupted teeth and assessing the extent of the anomaly.

Treatment approaches for Prikus Anamaias are tailored to the individual's specific clinical needs and the severity of the condition. Treatment goals typically include restoring proper masticatory function, improving aesthetics, and preventing further complications. Options may involve orthodontic intervention to facilitate tooth eruption or create space for prosthetic replacement of missing teeth. Surgical procedures might be necessary for the exposure and removal of impacted teeth or

for guided bone regeneration to enhance bone development in areas of missing teeth. Prosthodontic rehabilitation using partial or full dentures, fixed prostheses, or implants may be indicated to restore missing teeth and improve masticatory function. Interdisciplinary collaboration among dentists, orthodontists, oral and maxillofacial surgeons, and other specialists is often required to effectively manage this condition.

Materials and Methods

This study adopted a retrospective cohort design, analyzing data from pediatric patients (aged 6-12 years) diagnosed with Prikus Anamaias at [Name of Institution] over a 10-year period (2013-2023). Inclusion criteria comprised patients with confirmed Prikus Anamaias based on clinical examination (premature loss of deciduous teeth, failure of permanent tooth eruption) and radiographic assessment (panoramic and periapical radiographs) revealing impacted or missing permanent teeth. Patients with associated syndromes known to cause tooth agenesis (e.g., ectodermal dysplasia) were included separately for analysis.

REFERENCES

- 1.Rasulova D., Rasulova M. A CLINICAL CASE OF NEURODEGENERATIVE DISEASE OF THE TYPE OF LEWY BODY DISEASE WITH SEVERE COGNITIVE, AUTONOMIC DISORDERS //JOURNAL OF EDUCATION AND SCIENTIFIC MEDICINE. – 2023. – T. 2. – №. 1. – C. 45-49.
- 2.Bakhtiyarovna R. M. et al. REHABILITATION MEASURES IN PATIENTS WITH ISCHEMIC STROKE //INTERDISCIPLINE INNOVATION AND SCIENTIFIC RESEARCH CONFERENCE. – 2024. – T. 2. – №. 20. – C. 268-270.
- 3.Rasulova D. K., Rakhimbaeva G. S., Rasulova M. B. CLINICAL CASE OF REHABILITATION OF CENTRAL POST-STROKE NEUROPATHIC

PAIN-DEJERINE ROUSSY SYNDROME //British Medical Journal. – 2023. – T. 3. – №. 1.

4.Rasulova D. et al. Dynamics of restoring motor functions in post-stroke. – 2023.